Clim. Past, 8, 1091, 2012 www.clim-past.net/8/1091/2012/ doi:10.5194/cp-8-1091-2012 © Author(s) 2012. CC Attribution 3.0 License.





Corrigendum to

"The oxygen isotopic composition of phytolith assemblages from tropical rainforest soil tops (Queensland, Australia): validation of a new paleoenvironmental tool" published in Clim. Past, 8, 307–324, 2012

A. Alexandre¹, J. Crespin¹, F. Sylvestre¹, C. Sonzogni¹, and D. W. Hilbert^{1,2}

¹CEREGE, UMR7330, Aix-Marseille Université, CNRS, IRD, Europôle de l'Arbois, BP80, 13545 Aix en Provence, Cedex 04, France

²CSIRO Ecosystem Sciences, Tropical Forest Research Centre, P.O. Box 780, Atherton, QLD 4883, Australia

Correspondence to: A. Alexandre (alexandre@cerege.fr)

The sentence at the end of column 1 and beginning of column 2, p. 319, is unfortunately incomplete. One must read:

"The coefficient is in the range obtained for fossil diatoms (from -0.28 to $-0.49 \ ^{\circ}C^{-1}$, Juillet-Leclerc and Labeyrie, 1987; Shemesh et al., 1992), slightly larger than the value experimentally obtained for high temperature quartz (-0.24 to $-0.27 \ ^{\circ}{\rm o}^{\rm C^{-1}}$; Clayton et al., 1972; Matsuhisa et al., 1979), and larger than the value obtained for fresh diatoms (-0.19 to $-0.28 \ ^{\circ}{\rm C^{-1}}$, Brandriss et al., 1998; Moshen et al., 2005; Dodd and Sharp, 2010; Crespin et al., 2010)."

instead of:

"The coefficient is in the range obtained for fossil diatoms (from -0.28 to $-0.49 \ ^{\circ}C^{-1}$, Juillet-Leclerc and Labeyrie, 1987; Shemesh et al., 1992), slightly larger than the value experimentally obtained for high temperature quartz (-0.24 to $-0.27 \ ^{\circ}{\%} \ ^{\circ}C^{-1}$; Clayton et al., 1972; Matsuhisa et al., 1979), an $\ ^{\circ}{\%} \ ^{\circ}C^{-1}$, Brandriss et al., 1998; Moshen et al., 2005; Dodd and Sharp, 2010; Crespin et al., 2010)."